



9^ο ΣΥΜΠΟΣΙΟ ΟΜΑΔΩΝ ΕΡΓΑΣΙΑΣ


Της Ελληνικής Εταιρείας Χειρουργών
Θώρακος, Καρδιάς και Αγγείων

Η χειρουργική θεραπεία στην καρδιακή ανεπάρκεια ισχαιμικής αιτιολογίας – Από τη STICH στη STICHES

Πολυχρόνης Αντωνίτσης

Επ. Καθηγητής Καρδιοχειρουργικής Α.Π.Θ.





The NEW ENGLAND JOURNAL *of* MEDICINE

ESTABLISHED IN 1812

APRIL 23, 2009

VOL. 360 NO. 17

Coronary Bypass Surgery with or without Surgical Ventricular Reconstruction


Robert H. Jones, M.D., Eric J. Velazquez, M.D., Robert E. Michler, M.D., George Sopko, M.D., Jae K. Oh, M.D.,
Christopher M. O'Connor, M.D., James A. Hill, M.D., Lorenzo Menicanti, M.D., Zygmunt Sadowski, M.D.,
Patrice Desvigne-Nickens, M.D., Jean-Lucien Rouleau, M.D., and Kerry L. Lee, Ph.D.,
for the STICH Hypothesis 2 Investigators*



Υπόθεση 2




A Death from Any Cause or Hospitalization for Cardiac Causes



No. at Risk

CABG	499	319	270	220	99	23
CABG plus SVR	501	319	275	216	111	23

B Death from Any Cause



No. at Risk

CABG	499	434	417	363	201	59
CABG plus SVR	501	429	404	352	193	53

Jones RH, et al. New Eng J Med 2009.




Table 2. Outcomes and Subsequent Procedures.*


Variable	CABG Alone (N = 499) <i>number (percent)</i>	CABG with Surgical Ventricular Reconstruction (N = 501)	Hazard Ratio (95% CI)	P Value
Outcome				
Death from any cause or hospitalization for cardiac causes	292 (59)	289 (58)	0.99 (0.84–1.17)	0.90
Death from any cause	141 (28)	138 (28)	1.00 (0.79–1.26)	0.98
Death from any cause within 30 days after randomization (intention-to-treat analysis)	22 (4)	30 (6)		0.26
Death from any cause within 30 days after surgery†				
Intention-to-treat analysis	25 (5)	26 (5)		0.88
As-treated analysis	23 (5)	28 (6)		0.40
Hospitalization				
For any cause	272 (55)	268 (53)	0.98 (0.83–1.16)	0.82
For cardiac causes	211 (42)	204 (41)	0.97 (0.80–1.18)	0.73
Acute myocardial infarction	22 (4)	20 (4)	1.01 (0.54–1.87)	0.96
Stroke	31 (6)	23 (5)	0.77 (0.45–1.32)	0.35
Subsequent procedure‡				
CABG	0	1 (<1)		
Placement of left ventricular assist device	2 (<1)	2 (<1)		
Heart transplantation	2 (<1)	7 (1)		
Percutaneous coronary intervention	32 (6)	17 (3)		
Placement of pacemaker				
For resynchronization	31 (6)	30 (6)		
For heart rate	44 (9)	47 (9)		
Placement of implantable cardioverter–defibrillator	100 (20)	86 (17)		

Jones RH, et al. New Eng J Med 2009.


A CCS Angina Class



B NYHA Heart Failure Class



Jones RH, et al. New Eng J Med 2009.



Recommendations on revascularizations in patients with chronic heart failure and systolic LV dysfunction (ejection fraction $\leq 35\%$)

Recommendations	Class ^a	Level ^b	Ref ^c
CABG is recommended for patients with significant LM stenosis and LM equivalent with proximal stenosis of both LAD and LCx arteries.	I	C	
CABG is recommended for patients with significant LAD artery stenosis and multivessel disease to reduce death and hospitalization for cardiovascular causes.	I	B	112,288
LV aneurysmectomy during CABG should be considered in patients with a large LV aneurysm, if there is a risk of rupture, large thrombus formation or the aneurysm is the origin of arrhythmias.	IIa	C	
Myocardial revascularization should be considered in the presence of viable myocardium.	IIa	B	55
CABG with surgical ventricular restoration may be considered in patients with scarred LAD territory, especially if a post-operative LVESV index $<70 \text{ ml/m}^2$ can be predictably achieved.	IIb	B	291–295
PCI may be considered if anatomy is suitable, in the presence of viable myocardium, and surgery is not indicated.	IIb	C	

2014 ESC/EACTS Guidelines on myocardial revascularization

CORRESPONDENCE



Surgical Ventricular Reconstruction

The STICH trial: Misguided conclusions

Gerald D. Buckberg, MD,^a and Constantine L. Athanasuleas, MD^b



European Journal of Heart Failure (2010) 12, 1024–1027
doi:10.1093/eurjhf/hfq147

EDITORIAL

The STICH trial unravelled

Gerald D. Buckberg^{1*}, Constantine L. Athanasuleas², Andrew S. Wechsler³,
Friedhelm Beyersdorf⁴, John V. Conte⁵, and John E. Strobeck⁶


«Παραλειπόμενα» STICH

- Χαλάρωση κριτηρίων εισαγωγής κατά τη διάρκεια της μελέτης
- Ο τελο-συστολικός όγκος της ΑΚ (LVESI > 60 ml/m²) έπαψε να αποτελεί κριτήριο εισόδου στη μελέτη
- EF ≤ 35% παρέμεινε ως μοναδικό κριτήριο
- Μετατροπή από μελέτη ασθενών με καρδιακή ανεπάρκεια ισχαιμικής αιτιολογίας σε μελέτη ασθενών με στεφανιαία νόσο

«Παραλειπόμενα» STICH


- Λιγότεροι από 50% των ασθενών ήταν κλάση NYHA III-IV
- 13% των ασθενών δεν είχαν προηγούμενο ΕΜ
- 26% των ασθενών δεν εμφάνιζαν στένωση στον LAD
- Δε χρησιμοποιήθηκε μελέτη βιωσιμότητας του μυοκαρδίου
- Μόνο 50% των ασθενών εμφάνιζε ακινητική ή δυσκινητική περιοχή στο μυοκάρδιο
- Σε 41% των ασθενών δεν τοποθετήθηκε patch


CHF Survival: Registry Data



Buckberg GD, J Thorac Cardiovasc Surg 2010


% ελάτωση του τελο-συστολικού όγκου της ΑΚ (LVESVI)





A

White HD, et al. Circulation 1987;76:44-51.




Jones RH, et al. New Eng J Med 2009.

EUROPEAN
SOCIETY OF
CARDIOLOGY®


End-systolic volume following surgical ventricular reconstruction impacts survival in patients with ischaemic dilated cardiomyopathy

Ρόλος του LVESVI στην SVR




	0	12	24	36	48	60	72	84
≥60 mL/m ²	71	65	55	36	27	20	11	6
<60 mL/m ²	145	132	103	78	56	32	12	7

Διατοιχωματική ουλή



Scar_{Thick}




Wall_{Thick}



Transmurality (%): $\text{Scar}_{\text{Thick}} / \text{Wall}_{\text{Thick}} \times 100$ or $A/B \times 100$

e.g. $9\text{mm}/11\text{mm} \times 100 = 81.81\%$


Διατοιχωματική ουλή και βιωσιμότητα




Πάχος τοιχώματος

+


=




Βιωσιμότητα



Bull's-eye




Γεωμετρική αναδιαμόρφωση ΑΚ



Πριν

EDVI: 361 ml
EDVI: 154 ml/m²
ESV: 289 ml
ESVI: 123,5 ml/m²
EF: 20%
SVI: 20 ml/m²

Γεωμετρική αναδιαμόρφωση ΑΚ




Μετά SVR


EDVI: 60ml/m²

ESVI: 40 ml/m²

EF: 45%

SVI=31ml/m²






ORIGINAL ARTICLE

Coronary-Artery Bypass Surgery in Patients with Left Ventricular Dysfunction

Eric J. Velazquez, M.D., Kerry L. Lee, Ph.D., Marek A. Deja, M.D., Ph.D.,
Anil Jain, M.D., George Sopko, M.D., M.P.H., Andrey Marchenko, M.D., Ph.D.,
Imtiaz S. Ali, M.D., Gerald Pohost, M.D., Sinisa Gradinac, M.D., Ph.D.,
William T. Abraham, M.D., Michael Yui, M.S., F.R.C.S., F.R.A.C.S.,
Dorairaj Prabhakaran, M.D., D.M., Hanna Szwed, M.D., Paolo Ferrazzi, M.D.,
Mark C. Petrie, M.D., Christopher M. O'Connor, M.D.,
Pradit Panchavinnin, M.D., Lilin She, Ph.D., Robert O. Bonow, M.D.,
Gena Roush Rankin, M.P.H., R.D., Robert H. Jones, M.D.,
and Jean-Lucien Rouleau, M.D., for the STICH Investigators*







No. at Risk

Medical therapy	602	532	487	435	312	154	80
CABG	610	532	486	459	340	174	91


Velazquez et al., N Eng J Med, 2011



Velazquez et al., N Eng J Med, 2011


VOL. 374 NO. 16

Death from Any Cause (Primary Outcome)



Velazquez et al., N Eng J Med, 2016


Death from Any Cause or Cardiovascular Hospitalization



Velazquez et al., N Eng J Med, 2016


Ten-Year Outcomes After Coronary Artery Bypass Grafting According to Age in Patients With Heart Failure and Left Ventricular Systolic Dysfunction

An Analysis of the Extended Follow-Up of the STICH Trial (Surgical Treatment for Ischemic Heart Failure)




Petrie et al., Circulation, 2016

Καρδιακή ανεπάρκεια ισχαιμικής αιτιολογίας



Surgical Options in Advanced Heart Failure







Waveform Review

Weight(kg):

Bed No.:2

Patient ID:

Date of Birth:

Gender:

Patient Type:Adult


Admit Date:

Physician:


time:11/9/2016 3:55:16 PM Waveform end time:11/9/2016 3:55:27 PM Sweep Speed:25 mm/s 1 s/div



2

2




CABGX3,
Ηλεκτρική θύελλα
2-3 MTX > 40 απινιδώσεις στη ΜΕΘ





Surgical Options in Advanced Heart Failure



A Randomized Controlled Trial of Preoperative Intra-Aortic Balloon Pump in Coronary Patients With Poor Left Ventricular Function Undergoing Coronary Artery Bypass Surgery*

Marco Ranucci, MD, FESC¹; Serenella Castelvechio, MD, FESC¹; Andrea Biondi, MD²; Carlo de Vincentiis, MD²; Andrea Ballotta, MD, FESC¹; Alessandro Varrica, MD²; Alessandro Frigiola, MD²; Lorenzo Menicanti, MD²; for the Surgical and Clinical Outcome Research (SCORE) Group

TABLE 2. Primary Outcome (With Its Components) and Operative Mortality in the Two Groups

Outcome	Intra-Aortic Balloon Pump Group (n = 55)	Control Group (n = 55)	Odds Ratio (95% CI)	p
Surgical revision, n (%)	6 (11)	2 (3.6)	3.24 (0.62–18.84)	0.161
Prolonged mechanical ventilation, n (%)	17 (31)	15 (27)	1.19 (0.52–2.72)	0.675
Acute kidney injury, n (%)	5 (12)	2 (4.9)	2.71 (0.49–14.84)	0.251
Stroke, n (%)	1 (1.8)	0 (0)	Not applicable	0.315
Mediastinitis, n (%)	1 (1.8)	3 (5.5)	0.32 (0.03–3.19)	0.332
Major morbidity, n (%)	22 (40)	17 (31)	1.49 (0.68–3.33)	0.320
Operative mortality, n (%)	4 (7.3)	8 (14)	0.46 (0.13–1.43)	0.230

Ranucci et al., Crit Care Med, 2013

Effectiveness of prophylactic levosimendan in patients with impaired left ventricular function undergoing coronary artery bypass grafting: a randomized pilot study

Kyriakos Anastasiadis, Polychronis Antonitsis*, Konstantinos Vranis, Athanassios Kleontas, Christos Asteriou, Vassilios Grosomanidis, Paschalis Tossios and Helena Argiriadou








Figure 2: Time point recording of haemodynamic parameters.

Anastasiadis et al., ICVTS, 2016







Anastasiadis et al., ICVTS, 2016

Ισχαιμική μυοκαρδιοπάθεια

Στεφανιαία νόσος- Ισχαιμική
μυοκαρδιοπάθεια
(LVEF $\leq$ 40%)



CABG + IM εμφύτευση stem cells




Αυτόλογ
α


Ετερόλογ
α

AUTOLOGOUS BONE MARROW INJECTATE (im)
pluripotent population of stem cells (10^6)

- CD34+
- CD133+
- CD105+



Myocardial reperfusion at the infarcted and peri-infarct milieu.






Replacement of scarred tissue and regeneration of viable myocardium remains a challenging target of cell transplantation therapy.

ORIGINAL ARTICLE


Implantation of a Novel Allogeneic Mesenchymal Precursor Cell Type in Patients with Ischemic Cardiomyopathy Undergoing Coronary Artery Bypass Grafting: an Open Label Phase IIa Trial

Kyriakos Anastasiadis¹ • Polychronis Antonitsis¹ • Stephen Westaby² • Ajan Reginald³ • Sabena Sultan² • Argirios Doumas⁴ • George Efthimiadis⁵ • Martin John Evans⁶






Anastasiadis et al., JCTR, 2016






50 ετών


Πρωτοδιαγνωσθείσα
καρδιακή ανεπάρκεια
ισχαιμικής αιτιολογίας






		Endovolume	Blood volume
ED volume	:	389.8	324.5 ml
ES volume	:	328.3	259.0 ml
Ejection fraction	:	15.8	20.2 %
Stroke volume	:	61.5	65.6 ml
Stroke index	:	33.0	35.1 ml/beat/m ²
Cardiac output	:	3.6	3.9 l/min
Cardiac index	:	1.9	2.1 l/min/m ²

137092





- Βέλτιστη
φαρμακευτική αγωγή
- PCI
- Ενδεχόμενη μηχανική
υποστήριξη ως
θεραπεία προορισμού

